arXiv:hep-ph/0510229v1 18 Oct 2005

A role of parton charges and masses in the exclusive photon-photon production of meson pairs

A. V. Berezhnoy[1,*] and A. K. Likhoded[2,†]

[1]*SINP MSU, Moscow, Russia*

[2]*IHEP, Protvino, Russia*

The exclusive photon-photon production of the K-meson pairs has been analyzed within the partonic model of QCD for the kinematical conditions of BELLE experiment. The cross section dependences on partonic masses, charges, and a shape of K-meson wave function have been studied for the process under discussion.

1. INTRODUCTION

Not long ago the angular distributions for the process $\gamma\gamma \to K^+K^-$ have been measured by BELLE collaboration at energies 2.4-4.1 GeV [1]. The obtained results are in satisfactory agreement with the predictions of article [2], where the amplitude of exclusive production of meson pairs had been performed as the amplitude T_H of hard production of four partons convoluted with the parton amplitude distributions $\phi(x, Q)$ of the final mesons (x is a momentum fraction carried by valence parton and Q is a interaction scale):

$$A = \iint dx\, dy\, \phi^*(x, Q)\phi^*(y, Q)T_H. \tag{1}$$

The amplitude T_H of a hard subprocess of four massless quark production $\gamma\gamma \to q\bar{q}q\bar{q}$ was calculated in the frame work of perturbative QCD. It was rather difficult task because one need to take into account 24 Feynman diagrams of $O(\alpha^2\alpha_s^2)$.

The following parameterizations were used for $\phi(x, Q)$:

$$\phi(x, Q) \sim x(1 - x), \tag{2}$$

$$\phi(x, Q) \sim [x(1 - x)]^{\frac{1}{4}}, \tag{3}$$

$$\phi(x, Q) \sim \delta\left(x - \frac{1}{2}\right). \tag{4}$$

---

[*] Electronic address: `aber@ttk.ru`

[†] Electronic address: `likhoded@mx.ihep.su`

Transverse momenta of parton inside mesons, as well as partonic masses were neglected.

Now-days the wave function parametrization (2) is the most used for the study of processes with light mesons within the light-cone thechnique. The parametrization (3) was used in work [2], as an example of a function with a large partonic momentum dispersion. On the contrary, the wave function (4) describes a meson with fixed momenta of constituent partons. The latter parametrization has been used many times to study the processes with heavy quarkonia (see for example a set of papers devoted to B_c-production in different interactions [3]). Also the parameterization (3) has been used by us to estimate the production cross sections and the charge asymmetries in the charm meson photoproduction [4].

One of the aims of this study is to estimate the role of partonic masses in the inclusive photonic production of meson pairs. For this purpose we have taken into account the mass corrections in the hard process amplitude T_H.

The other aim of the research is to note one more time, that the values of partonic charges influence strongly the shapes of differential distributions. Our calculations, as well as the results of [2] are indicative of the cardinal difference between the angle distributions for charged meson pairs and ones for neutral meson pairs. This difference is clearly seen from the formula for the cross section distribution over cosine of azimuthal angle Θ in c.m.s. for the production of pseudoscalar meson pairs, which had been obtained in [2]:

$$\frac{d\sigma}{d\cos\Theta} \sim \frac{(e_1-e_2)^4}{\sin\Theta^4} + \frac{2e_1e_2(e_1-e_2)^2}{\sin\Theta^2} g[\Theta,\phi] + 2(2e_1e_2)^2 g^2[\Theta,\phi], \tag{5}$$

where $g[\Theta,\phi]$ is a function of angle and of $\phi(x)$, e_1 and e_2 are the quark charges (the charges of produced mesons are equal to $\pm(e_1-e_2)$). It is worth to underline that the both terms in (5) can contribute essentially to K^+K^--pair production process. That is why the approximation to the cross section in the form of $1/\sin\Theta^4$, which was used in [1], does not take into account all features of the model under discussion. For example, for the δ-shaped amplitude $\phi(x)$ and for massless partons $g[\Theta,\phi] \equiv 1$ and the cross section of K^+K^--pair depends on Θ as follows:

$$\frac{d\sigma}{d\cos\Theta} \sim \frac{1}{\sin\Theta^4} - \frac{4}{9\sin\Theta^2}. \tag{6}$$

Let us to note that the term proportional to $1/\sin^2\Theta$ allows to slightly improve the description of experimental data published in [1].

As it was already mentioned above, the model under discussion predicts the essential difference between the cross section distributions for neutral meson pair production and for charged meson pair production. It is especially interesting because the other models [5] does not predict an essential dependence of the differential cross section on parton charges.

The third aim of the present work is to find-out the influence of a shape of $\phi(x, Q)$ on the behavior of cross section distributions for massive partons, as well as massless ones.

2. AN AMPLITUDE DISTRIBUTION AND A MASS OF A SYSTEM CONSTITUTED FROM TWO PARTONS

Before to state the main results of the study let us return to the discussion of amplitude distribution shape to make two remarks about this. As it is mentioned above, the approximation (2) is the most used for the case of light mesons constituted from massless partons. The validity of parametrization (2) for calculations is widely discussed in the literature [6], and we do not consider here this question. Our observation concerns the other aspect of the problem. Let us suppose that the valence quarks q_1 and q_2 have the effective masses m_{q_1} and m_{q_2}. Let us also suppose that these partons carry practically the total momentum of meson. Then neglecting their transverse momenta we obtain a following expression for the mass squared of such two parton system:

$$m^2(x) = \left(\sqrt{x^2p^2 + m_{q_1}^2} + \sqrt{(1-x)^2p^2 + m_{q_2}^2}\right)^2 - p^2, \tag{7}$$

where x is a fraction of meson momentum p carried away by the parton q_1. At large momenta $m^2(x)$ can be represented as

$$m^2(x)\Big|_{p\to\infty} = \frac{m_{q_1}^2}{x} + \frac{m_{q_2}^2}{1-x}. \tag{8}$$

It is easy to show that the function $m(x)$ reaches the minimum value $m_{q_1} + m_{q_2}$ at $x = m_{q_1}/(m_{q_1} + m_{q_2})$.

If $m_{q_1} = m_{q_2}$ then $m^2(x) \sim 1/(x(1-x))$ and the amplitude (2) depends on x as $1/m^2(x)$. Thus, one could suppose that for unequal masses m_{q_1} and m_{q_2} a following heuristic expression for the parton amplitude can be used:

$$\phi(x) \sim \frac{1}{m^2(x)} \sim \left(\frac{r^2}{x} + \frac{1}{1-x}\right)^{-1}, \tag{9}$$

where $r = m_{q_1}/m_{q_2}$. At least, for the reasonable value of parameter $r = 2$ the received distribution is practically indistinguishable from one represented in work [7] for K-meson.

The other remark is that even for a heavy quarkonium the value of $m(x)$ can not be identify with the quarkonium mass because this mass can not depend on x. Nevertheless, we think that a physical meaning have only the values of x, at which the difference between $m(x)$ and the meson mass M not larger than a coupling energy value Λ. If the both quarks are heavy then $M \approx m_{q_1} + m_{q_2}$, $M \gg \Lambda$, and we arrive at the following inequality:

$$m(x) - m_{q_1} - m_{q_2} \lesssim \Lambda. \tag{10}$$

In addition to that, if $m_{q_1} = m_{q_2}$, then from (8) and (10) we obtain:

$$\frac{1}{2} - \frac{1}{2}\sqrt{\frac{\Lambda}{m_q}} \lesssim x \lesssim \frac{1}{2} + \frac{1}{2}\sqrt{\frac{\Lambda}{m_q}}. \tag{11}$$

Using (11) one can easy to estimate a region of reasonable value of x both for charmonium ($0.3 \lesssim x \lesssim 0.7$) and bottomium ($0.35 \lesssim x \lesssim 0.65$).

Thus, following [2], one can essentially simplify the problem if suppose that the amplitude of exclusive production of meson pair can be subdivided into the hard amplitude of parton production and the soft adronization amplitude. But such a factorization cause the vagueness of meson mass. Different values of x correspond to different values of a mass of system with two partons. In other words, we replace a meson with a fixed mass value to a wave package constituted by partons. Such an object can not be characterized by the definite mass value. Nevertheless, a rather wide region of x variable exits where the mass of two parton reach physically reasonable values as for light mesons, as for heavy ones. Therefore the factorization proposed in [2] is physically justified.

3. CALCULATION RESULTS

In the present work we do not concern the problem of absolute normalization of the cross section and discuss only the shape of cross section distribution with respect to $\cos\Theta$ depending on the following factors:

1. a value of meson charge,
2. values of effective parton masses,
3. a shape of parton amplitude $\phi(x)$.

At the present time there are many work [6] devoted to study of different parameterizations of parton amplitude and their behavior. Nevertheless, as it was experienced by previous calculations, the shape of parton amplitude can not be reconstructed in details from the behavior of cross section or decay width. The process under discussion is not an exception and the differential cross section of the meson pair production allows us to determine only qualitative characteristics of the parton amplitude $\phi(x)$. That is why use the rough model represented in [2] to obtain our predictions.

The cross section calculations for K-meson production have been done within the following assumption sets about a parton amplitude shape and parton masses:

1. s-quark mass equals to $m_s = 0.2$ GeV, the masses of u- and d-quarks equal to $m_q = 0.1$ GeV and the momentum fractions carried by the partons are fixed: $\phi(x) \sim \delta(x - m_s/(m_s + m_q))$.
2. $m_s = 0.2$ GeV, $m_q = 0.1$ GeV, and the parton amplitude are spread as much as possible: $\phi(x) = const > 0$ at $0.25 < x < 0.95$ and $\phi(x) = 0$ at other x. The region of x at which the parton amplitude reaches nonzero values has been estimated from the condition (10).
3. The parton masses are negligibly small, and $\phi(x) \sim \delta(x - 2/3)$.
4. The parton masses are negligibly small, and $\phi(x)$ have a shape (9).

Of course these extreme variants are not realized in the nature. However the study of the itemized assumption sets allows to better understand a new experimental data.

Fig. 1a presents the normalized differential cross sections of pair production of charged K-mesons with respect to $|\cos\Theta|$ for the process $\gamma\gamma \to K^+K^-$ at 2.65 GeV for the supposition sets under discussion: (1) (solid curve), (2) (dashed curve), (3) (dotted curve), and (4) (dashed-dotted curve). The experimental data of Belle Collaboration are also given in Fig. 1a for comparison. All four sets describe data fairly well. Surprisingly, that the most fair description has been achieved for the variant (3) which can be approximated by the formula (6) with a high accuracy. However one can not come to a final decision.

Fig. 1b presents the differential cross sections of pair production of charged vector and pseudoscalar K-mesons ($\gamma\gamma \to K^+K^{*-}(K^{*+}K^-)$) normalized to the cross section value of pair production of charged pseudoscalar mesons. It is seen from Fig. 1b that the cross section of the process under discussion depends on Θ as $\sim \cos^2\Theta/\sin^4\Theta$. Also one can see that the relative yield of pseudoscalar-vector pairs for the "spread" parton amplitudes is essentially smaller than for the δ-shaped one.

As it is seen from Fig. 1c for the case of massless partons the yield of the vector-vector pairs is smaller than for the case of massive ones. On comparing Fig. 1c, 1a, and 1b, it is clear, that the cross section of the process $\gamma\gamma \to K^{*+}K^{*-}$ depends on $|\cos\Theta|$ weaker than the cross sections of the processes $\gamma\gamma \to K^+K^-$ and $\gamma\gamma \to K^+K^{*-}(K^{*+}K^-)$.

The differential production cross sections of the pairs of neutral pseudoscalar K-mesons are crucially differ from the cross sections obtained for the charged pair production. The cross section for process $\gamma\gamma \to K^+K^-$ strongly depends on $|\cos\Theta|$. On the contrary, the distribution over $|\cos\Theta|$ for the process $\gamma\gamma \to K^0\bar{K}^0$ is practically flat for the δ-shaped parton amplitude and slightly curved for the spread parton amplitude (see Fig. 1d).

It is clear from Fig. 4e, that the process $\gamma\gamma \to K^0\bar{K}^{*0}(K^{*0}\bar{K}^0)$ is strongly suppressed in the case of massless partons. For the scheme (1) with massive partons and fixed parton momenta the cross section distribution over $|\cos\Theta|$ with fair accuracy can be approximated by the function $|1/\sin^3\Theta|$. A spreading of parton amplitude within scheme (2) leads to the increase of the relative $K^0\bar{K}^{*0}$-pair yield and to the slight alteration of the distribution shape.

The values of parton charges influence strongly the behavior of cross section distribution of pseudoscalar-pseudoscalar and pseudoscalar-vector pairs, whereas the distribution shape of vector-vector pair production practically does not depend on parton charges, as it is clear from comparing Fig. 1f and Fig. 1c were the cross section distributions over $|\cos\Theta|$ are shown the processes $\gamma\gamma \to K^{*0}\bar{K}^{*0}$ and $\gamma\gamma \to K^{*+}K^{*-}$, correspondingly.

Fig. 2 presents the same distributions as Fig. 1 but at the photon-photon interaction energy 3.15 GeV. One can see that there is no essential difference between the distributions at 2.65 GeV (Fig. 1) and at 3.15 GeV (Fig. 2).

4. CONCLUSIONS

Our calculations show that the shapes of differential cross section of the pseudoscalar meson pair production in the photon-photon interaction is determined in general by the meson charge. For the process $\gamma\gamma \to K^+K^-$ the distribution over $\cos\Theta$ have a strongly marked peripheral character, whereas the analogous distribution for the process $\gamma\gamma \to K^0\bar{K}^0$ is practically flat. It is worth to note that the partonic mass values and the shape of parton amplitude influence weakly the differential cross section behaviour.

The differential cross section shapes of pseudoscalar-vector production are determined by the meson charge too. But the relative yield of pseudoscalar-vector pairs and vector-vector pairs depends essentially on the parton amplitude parameterization, as well as on the parton mass values. It is interesting that the prediction of cross section value for the process $\gamma\gamma \to K^0\bar{K}^{*0}(K^{*0}\bar{K}^0)$ can be essentially reduced by using a spread parton amplitude, whereas the cross section value for the process $\gamma\gamma \to K^+K^{*-}(K^{*+}\bar{K}^-)$ is more sensitive to the choice of parton masses values and shape of parton amplitude. For the massless partons the cross section of the latter process is negligibly small independently on the chosen parton amplitude parametrization.

The vector-vector pair production have a weakly marked peripheral character. The relative yield of vector-vector pair, as well as the shape of cross section distributions weakly depend on suppositions about both parton distributions and parton mass values.

The neutral vector-vector pair production have a marked peripheral character for spread parton amplitudes and is practically isotropic for the δ-shaped amplitude.

To conclude, the present research shows that all tree discussed factors (the meson charge value, the shape of parton amplitude, and the values of parton masses) can essentially influence the exclusive meson pair production in the photon-photon interaction. Nevertheless, the role of meson charge value must be especially stressed, because a change of meson charge leads to the most crucial change in the meson production. That is why it should be very interesting to compare the experimental results on the pair production of charged K-mesons [1] with experimental results on the pair production of neutral K-mesons.

This work was supported in part by the Russian Foundation for Basic Research (project no. 04-02-17530) and CRDF (grant no. M0-011-0). It was also funded within the Program for Support of Leading Scientific Schools (grant no. 1303.2003.2).

---

1. H. Nakazawa *et al.* [Belle Collaboration], hep-ph/0412058.
2. S. J. Brodsky and G. P. Lepage, Phys. Rev. D **24**, 1808(1981).
3. A. V. Berezhnoi, A. K. Likhoded, and M. V. Shevlyagin, Phys. Atom. Nucl. **58**, 672 (1995); A. V. Berezhnoi, A.K. Likhoded, and O. P. Yushchenko, Phys. Atom. Nucl. **59**, 709 (1996); A. V. Berezhnoi, V. V. Kiselev, and A. K. Likhoded, Phys. Atom. Nucl. **60**, 100 (1997); A. V. Berezhnoy, V. V. Kiselev, and A. K. Likhoded, Z. Phys. A **356**, 79 (1996); A.V. Berezhnoi, V.V. Kiselev, A.K. Likhoded, and A.I. Onishchenko, Phys. Atom. Nucl. **60**, 1729 (1997).
4. A. V. Berezhnoy, V. V. Kiselev, and A. K. Likhoded, Phys. Atom. Nucl. **63**, 1595(2000); A. V. Berezhnoy, V. V. Kiselev, and A. K. Likhoded, Phys. Rev. D **62**, 074013 (2000); A. V. Berezhnoy, V. V. Kiselev, and A. K. Likhoded, Phys. Atom. Nucl. **66**, 769 (2003).
5. M. Diehl, P. Kroll and C. Vogt, Phys. Lett. B **532**, 99 (2002).
6. V. L. Chernyak and I. R. Zhitnitsky, Nucl. Phys. B **246**, 52 (1984); A. R. Zhitnitsky, I. R. Zhitnitsky and V. L. Chernyak, Sov. J. Nucl. Phys. **38**, 645 (1983); P. Ball and V. M. Braun, Phys. Rev. D **54**,

2182 (1996); P. Ball and V. M. Braun, NORDITA-98-51-HE, Apr. 1998; Talk given at 3rd Workshop on Continuous Advances in QCD (QCD 98), Minneapolis, 16—19 Apr. 1998, p. 125.

7. P. Ball, V. M. Braun, Y. Koike and K. Tanaka, Nucl. Phys. B **529**, 323 (1998); P. V. Chliapnikov, V. G. Kartvelishvili, V. V. Knyazev and A. K. Likhoded, Nucl. Phys. B **148**, 400 (1979).



Figure 1. The differential cross sections of K-meson pair production in the photon-photon interaction at 2.65 GeV with respect to $|\cos\Theta|$ for the assumption sets under discussion: (1) $m_s = 0.2$ GeV, $m_u = m_d = 0.1$ GeV, $\phi(x) \sim \delta(x - m_s/(m_s + m_q))$ (solid curve); (2) $m_s = 0.2$ GeV, $m_u = m_d = 0.1$ GeV, $\phi(x) = const > 0$ at $0.25 < x < 0.95$ and $\phi(x) = 0$ at other x (dashed curve); (3) the parton masses are negligibly small, $\phi(x) \sim \delta(x - 2/3)$ (dotted curve); (4) the parton masses are negligibly small, $\phi(x)$ have a shape (9) (dashed-dotted curve). See the text for the detailed explanation.



Figure 2. The same as in Fig. 1, but for the interaction energy 3.15 GeV.